arXiv:1211.3481v2 [cond-mat.str-el] 19 Nov 2012

Realization of 2-channel Kondo effect in a junction of three quantum Ising chains

A. M. Tsvelik[1]

[1]*Department of Condensed Matter Physics and Materials Science,*
Brookhaven National Laboratory, Upton, NY 11973-5000, USA

(Dated: November 21, 2012)

It is shown that a junction of three quantum Ising chains (Y-junction) can be described as the 2-channel Kondo model in a box which size is of the order of the Ising model correlation length with spin S=1/2 localized at the junction. The local spin is composed of the zero energy boundary Majorana modes of the Ising models.

PACS numbers: 75.10.-b, 72.10.Fk

Majorana fermions in general and Majorana zero energy modes (MZEMs) in particular have become very popular topic. The interest is mainly caused by non-Abelian properties of MZEMs related to the fact that for a set of N modes the corresponding operators satisfy the Clifford algebra

$$\{\gamma_i, \gamma_j\} = \delta_{ij}, \quad i,j = 1,...N, \tag{1}$$

and as such represent a realization of a spinor representation of the O(N) group.

Theory predicts that MZEMs will appear at topological defects such as vortices in two- and domain walls in one dimension. Although in one dimension the task of generating Majorana modes is much easier, it is more difficult to take advantage of the non-Abelian nature of MZEMs since to exchange them one needs somehow to escape from 1D. It has been suggested that one can resolve the conundrum with a help of the so-called Y-junction. In [1] it was suggested to use Y-junctions (called T-junctions in that paper) to implement braiding operations of Majorana fermions.

In this paper I study a junction of three quantum Ising models coupled together at one point. The Hamiltonian is the one of three quantum Ising chains

$$H_{Ising,p} = \sum_{j=1}^{N}\Big[-J\sigma^z_p(j)\sigma^z_p(j+1) + h\sigma^x_p(j)\Big] \tag{2}$$

coupled together in the Y-junction:

$$H = -J_{12}\sigma^z_1(1)\sigma^z_2(1) - J_{23}\sigma^z_2(1)\sigma^z_3(1) - J_{13}\sigma^z_1(1)\sigma^z_3(1) + \sum_{p=1}^{3} H_{Ising,p} \tag{3}$$

I assume that $0 < J_{pq} << J$. Models (2) are equivalent to noninteracting Majorana fermions by means of Jordan-Wigner transformation and their boundaries contain MZEMs. It will be shown that for nearly critical chains $|J-h| << J$ the MZEMs located at the junctions are subject of intense screening by gapless bulk modes and model (3) is equivalent to the overscreened two-channel Kondo model. The latter model possesses non-Fermi liquid fixed point [2] with a residual ground state entropy [3],[4].

In thermodynamic limit $N \to \infty$ at zero temperature model (2) is in the ferromagnetically ordered state at $J > h$, where far from the boundary the magnetization is finite $\langle\sigma^z\rangle \sim (J-h)^{1/8}$; at $J < h$ the ground state is disordered (paramagnetic) $\langle\sigma^z\rangle = 0$. The point $J = h$ is critical. At $J \geq h$ the ground state is double degenerate in the thermodynamic limit. This degeneracy is related to MZEMs located at the opposite boundaries of the system.

Majorana fermion a is a linear combination of fermionic creation and annihilation operator

$$a = (f + f^+)/2, \quad a^2 = 1/2, \quad a = a^+, \tag{4}$$

and as such cannot be used to create or destroy any state. To do this one needs a pair of Majorana fermions a_1, a_2 from which creation and annihilation operators can be constructed:

$$f = a_1 + \mathrm{i}a_2, \quad f^+ = a_1 - \mathrm{i}a_2. \tag{5}$$

The zero modes located at different boundaries overlap; their tunneling Hamiltonian is

$$H_{tunn} = 2\mathrm{i}ta_1a_2 = t(f^+f - 1/2). \tag{6}$$

Thus for a finite chain the ground state degeneracy is broken, but since t decays fast with the system size, for sufficiently large systems one can think about boundary Majorana modes as independent. Due to their nonlocality such Majorana modes have attracted a particular attention in conjunction with quantum computation.

Let us return to model (2). As I have said, it is equivalent to the model of noninteracting Majorana fermions. In the continuum limit $|J-h| << J$ it is described by the Lagrangian

$$L = \int_{-\infty}^{\infty}\mathrm{d}x\Big[\frac{1}{2}\chi_R(\partial_\tau - \mathrm{i}v\partial_x\chi_R) + \frac{1}{2}\chi_L(\partial_\tau - \mathrm{i}v\partial_x\chi_L) + \mathrm{i}M(x)\chi_R\chi_L\Big] \tag{7}$$

where $v \sim J$ and $\chi_{L,R}$ are real anticommuting fields. Here I introduced a coordinate dependent mass term to simulate the effect of the boundaries. At $0 < x < L = Na_0$ we have $M(x) = h - J$, outside of this interval it interpolates to a large positive value $M \sim J$. If $h < J$ then the function $M(x)$ changes sign at the boundaries. Then the corresponding Dirac equation for the fermions has solutions with zero energy localized at $x = 0, x = L$ (I will treat these points as located infinitely far from each other and neglect all tunneling). These are Majorana zero modes (MZEMs). If $h > J$ (the disordered phase) $M(x)$ does not change sign and such solutions do not exist.

It is convenient to isolate the zero mode from other modes which was done in [6] where the following equivalent representation of the Lagrangian (7) was introduced (see Eq.(4.22) in [6]):

$$L = \int_0^\infty \mathrm{d}x\Big[\frac{1}{2}\chi_R(\partial_\tau - \mathrm{i}v\partial_x\chi_R) + \\ \frac{1}{2}\chi_L(\partial_\tau - \mathrm{i}v\partial_x\chi_L) + \mathrm{i}M\chi_R\chi_L\Big] + \frac{\mathrm{i}}{2}a\partial_\tau a, \\ (\chi_L - \chi_R)_0 = 0, \quad (M > 0), \tag{8}$$

In this representation the zero mode a is present explicitely and the bulk Lagragian is supplemented by the boundary condition. At time scales $|\tau| << |M|^{-1}$ the following formula is valid (see Eq.(4.25) in [6]) for p-th chain:

$$\sigma^z(1)_p = \mathrm{i}a_p(\chi_p^L + \chi_p^R)_0, \tag{9}$$

where a_p are local Majorana's living on the boundary. It has a dimension $[mass]^{1/2}$. This representation reproduces the known result for the correlation function of boundary spins:

$$\langle\sigma^z(\tau,1)\sigma^z(0,1)\rangle = \langle\hat{T}a(\tau)a(0)\rangle\langle\hat{T}\chi(\tau)\chi(0)\rangle = \\ \mathrm{sign}\tau \times \frac{1}{2\pi\tau} = \frac{1}{2\pi|\tau|}. \tag{10}$$

Substituting (9) into (3) and taking the continuum limit of the individual chains (7) we get the following effective Hamiltonian (we extended the space interval to the entire axis, but left only one chirality of the fermions: $\chi(x) = \theta(x)\chi_R(x) + \theta(-x)\chi_L(-x)$.):

$$H_{eff} = \mathrm{i}\sum_p G_p S^p \epsilon_{pqt}\chi_q(0)\chi_t(0) + \\ \frac{\mathrm{i}}{2}\int_{-\infty}^{\infty}\mathrm{d}x\chi_p\partial_x\chi_p, \tag{11}$$

where $G_1 = J_{23}, G_2 = J_{13}, G_3 = J_{12}$, and

$$S^p = \frac{\mathrm{i}}{2}\epsilon_{pqt}a_q a_t, \tag{12}$$

are components of spin S=1/2 [7] and the fermionic bilinears are $\mathrm{SU}_2(2)$ currents:

$$\mathcal{J}^a = \frac{\mathrm{i}}{2}\epsilon_{abc}\chi_b\chi_c.$$

The level of the current algebra is equal to the number of channels of this Kondo model. Model (11) describes the two-channel Kondo problem written in the form introduced in [5]. If the bare interactions are ferromagnetic ($-J_{ab} < 0$, the same sign as the bulk exchange), the Kondo exchange is antiferromagnetic and the interaction scales to the intermediate coupling critical point. Otherwise it is marginally irrelevant. A similar Kondo model (the 4-channel one) has recently been suggested in [8] in the context of the so-called topological Kondo effect. In [9] it was found that 4-channel Kondo model describes Y-junction of XX spin S=1/2 chains. This paper also contains an elegant derivation of the fermionic lattice Hamiltonian.

As was established in [5], the low energy Lagrangian describing the junction dynamics at energies less than the Kondo temperature T_K:

$$L_{eff} = \frac{\mathrm{i}}{2}\epsilon\partial_\tau\epsilon + g\epsilon\chi_1(0)\chi_2(0)\chi_3(0) + \sum_{p=1}^{3}L[\chi_p], \tag{13}$$

where $g \sim T_K$ and $L[\chi_p]$ describes three chiral Majorana modes of the bulk. The Kondo scale T_K is a function of the bare couplings and the ultraviolet cut-off J, for equal couplings $G_1 = G_2 = G_3$ it is exponentially small in G. The critical point is characterized by a single zero energy Majorana fermion coupled to the bulk by the irrelevant operator. Thus the Y-junction of critical Ising models yields one Majorana fermion located at the junction and weakly coupled to the environment as described in [5]. This conclusion holds irrespectively of the ratios between the couplings G_a; it is well known that the multichannel Kondo model fixed point is unaffected by the anisotropy [10]. This is a remarkable fact meaning that the low energy properties of the junction are robust with respect to anisotropy of the couplings. Another remarkable fact is that the local pseudospin (12) constructed from the MZEMs of different chains is not a local operator of Ising chains and therefore there is no "magnetic field" which can attached to it. Thus the 2-channel critical point cannot be destroyed by local perturbations. As far as nonlocal ones are concerned, the Kondo effect will not survive if at least on of the chains becomes noncritical. In the first approximation we can say that making the bulk massive is equivalent to putting our Kondo model in a box of size M^{-1}. As long the Kondo temperature exceeds M, not much changes. Otherwise, as we said, the Kondo effect disappears and the junction bahaves as S=1/2.

Let us consider two Y-junctions coupled together. Zero modes from one of them we denote η^-, from the other η^+. The Hamiltonian (2,3) acquires a correction in the form of tunneling term:

$$\mathrm{i}t_a\eta_a^+\eta_a^-, \quad t \sim 1/L. \tag{14}$$

This term competes with the Kondo screening which can be established looking at the mean field solution where the interaction is decoupled by fields Δ_σ so that the Lagrangian becomes (I consider the isotropic interaction):

$$L[\chi^\sigma] + \Delta_\sigma^2/2G + \mathrm{i}\Delta_\sigma\eta_a^\sigma\chi_a^\sigma(0) + \mathrm{i}t\eta_a^+\eta_a^-, \quad \sigma = \pm 1. \tag{15}$$

The saddle point action is

$$\frac{1}{2G}(\Delta_+^2 + \Delta_-^2) - \frac{3}{8\pi}\int \mathrm{d}\omega \ln[(|\omega| + \Delta_+)(|\omega| + \Delta_-) + t^2] \quad (16)$$

yielding

$$\Delta_\sigma = T_K - t, \quad (t < T_K); \quad \Delta_\sigma = 0, \quad (t > T_K) \qquad (17)$$

where $T_K \sim \exp(-Const/G)$. From here we may conclude that the Kondo effect disappears at $t > T_K$.

The current study indicates that the Y-junction of nearly critical quantum Ising models behaves as an active element where MZEMs undergo screening by the bulk excitations. At low energies the system scales to overscreened 2-channel Kondo model critical point. This is an interesting fact by itself since realization of overscreened Kondo effect is a notoriously difficult task. At the same time nearly critical quantum Ising systems can be realized in Josephson junction arrays [11].

I am grateful to A. A. Nersesyan for bringing this problem to my attention and extended subsequent discussions and to B. L. Altshuler and A. B. Zamolodchikov for the interest to the work. The work was supported by US DOE under contract number DE-AC02 -98 CH 10886.

---

[1] J. Alicea, Y. Oreg, G. Refael, F. von Oppen and M. P. A. Fisher, Nature Phys. **7**, 412 (2011).

[2] P. Nozieres and A. Blandin, J. Phys. **41**, 193 (1980).

[3] A. M. Tsvelick, P. B. Wiegmann, Z. für Phys. B**54**, 201 (1984); J. Stat. Phys. **38**, 125 (1985).

[4] A. M. Tsvelik, J. Phys. C**18**, 159 (1985).

[5] P. Coleman, L. B. Ioffe, A. M. Tsvelik, Phys. Rev. B**52**, 6611 (1995).

[6] S. Ghoshal and A. B. Zamolodchikov, Int. J. Mod. Phys. A**9**, 3841 (1994); ibid. p.4353.

[7] J. L. Martin, Proc. R. Soc.London Ser. A**251**, 536; R. Casalbuoni, Nuovo Cimento **33**, 389 (1976).

[8] B. Beri and N. R. Cooper, Phys. Rev. Lett. **109**, 156803 (2012).

[9] N. Crampe, A. Trombettoni, arXiv:1210.7143.

[10] I. Affleck, A. W. W. Ludwig, H. B. Pang, D. L. Cox, Phys. Rev. B **45**, 7918 (1992).

[11] M. Gershenson, private communication.